

ANALYTICITY FOR SINGULAR SUMS OF SQUARES OF DEGENERATE VECTOR FIELDS

DAVID S. TARTAKOFF

ABSTRACT. Recently, J.J. Kohn in [6] proved hypoellipticity for

$$(*_k) \quad P = L\bar{L} + \bar{L}|z|^{2k}L \quad \text{with} \quad L = \frac{\partial}{\partial z} + i\bar{z}\frac{\partial}{\partial t},$$

i.e.,

$$-P = \bar{L}^* \bar{L} + (\bar{z}^k L)^* \bar{z}^k L,$$

a singular sum of squares of complex vector fields on the complex Heisenberg group, an operator which exhibits a *loss* of $k-1$ derivatives. Subsequently, in [4], M. Derridj and D. S. Tartakoff proved analytic hypoellipticity for this operator using rather different methods going back to [8], [9]. Together with A. Bove and J. J. Kohn, Derridj and Tartakoff in [1] gave an alternative proof of the C^∞ hypoellipticity of $*_k$ in the style of [4]. In this note, we consider the equation

$$(*_{m,k}) \quad P = L_m \bar{L}_m + \bar{L}_m |z|^{2k} L_m \quad \text{with} \quad L_m = \frac{\partial}{\partial z} + i\bar{z}|z|^{2m} \frac{\partial}{\partial t},$$

for which the underlying manifold is only of finite type, and prove analytic hypoellipticity. This operator is subelliptic with large loss of derivatives but the exact loss is irrelevant for analytic hypoellipticity.

1. INTRODUCTION AND STATEMENT OF THEOREMS

J.J. Kohn's recent paper [6], inspired by work of Siu on singular metrics and the implied applications, studied a singular sum of squares of complex vector fields on the (complex) Heisenberg group:

$$(*_k) \quad P = -L\bar{L} - \bar{L}|z|^{2k}L \quad \text{with} \quad L = \frac{\partial}{\partial z} + i\bar{z}\frac{\partial}{\partial t},$$

i.e.,

$$P = \bar{L}^* \bar{L} + (\bar{z}^k L)^* \bar{z}^k L,$$

He showed that this operator was hypoelliptic but loses $k-1$ derivatives (in Sobolev norms), and a note by Christ showed that the addition of

Date: February 1, 2008.

the square of another vector field, $\frac{\partial^2}{\partial s^2}$ to P , destroyed hypoellipticity completely.

Subsequently, in [4], M. Derridj and D. S. Tartakoff proved analytic hypoellipticity for this operator using rather different methods going back to [8], [9]. In [1], together with A. Bove and J. J. Kohn, M. Derridj and D.S. Tartakoff then gave an alternative proof of the C^∞ hypoellipticity of $*_k$ in the style of [4].

In this paper, we consider the more degenerate equation

$$(*_{m,k}) \quad P = P_m = L_m \overline{L_m} + \overline{L_m} |z|^{2k} L_m \quad \text{with} \quad L_m = \frac{\partial}{\partial z} + i\bar{z}|z|^{2m} \frac{\partial}{\partial t},$$

for which the underlying manifold is only of finite type, and prove analytic hypoellipticity for this operator.

Theorem 1. *The operator P_m is analytic hypoelliptic: $P_m u \in C^\omega \rightarrow u \in C^\omega$.*

2. THE *a priori* ESTIMATE

As in [4] we work with $v \in C_0^\infty$ and denote by Λ the ‘pseudodifferential’ operator with symbol $\lambda(z, t; \zeta, \tau) = (1 + |\tau|^2)^{1/2}$. While Λ is not a true pseudodifferential operator, for ζ different from zero the operator P_m is elliptic, hence analytic hypoelliptic, near $z = 0$ and for z different from 0, the operator P_m has symplectic characteristic variety and satisfies maximal estimates in L and $\overline{L}$, hence is even analytic hypoelliptic there by [8], [10], [9]. Thus we will need only study the behavior of powers of $\partial/\partial t$ applied to the solution u locally. We shall do so in L^2 norms with appropriate powers of Λ .

The *a priori* estimate satisfied by P_m is actually better than that satisfied by P_0 but as is well known, analyticity does not use the precise degree of subellipticity in an important way. Thus we give a simple proof of the estimate given in Kohn’s paper [6].

Proposition 2.1. *For all values of m , we have*

$$\|\overline{L_m} v\|^2 + \|\bar{z}^k L v\|^2 + \|v\|_{-\frac{k-1}{2}}^2 \leq C |(P_m v, v)|, \quad v \in C_0^\infty.$$

For any $r < 0$, since Λ commutes with everything, we have:

$$\|\Lambda^r v\|^2 = ([L_m, z] \Lambda^r v, \Lambda^r v) = (z \Lambda^r v, \Lambda^r \overline{L_m} v) - (L_m \Lambda^{r-\frac{1}{2}} v, \bar{z} \Lambda^{r+\frac{1}{2}} v)$$

so

(2.1)

$$\|\Lambda^r v\|^2 \leq s.c. \|\Lambda^r v\|^2 + l.c. \|\overline{L_m} v\|^2 + s.c. \|L_m \Lambda^{r-\frac{1}{2}} v\|^2 + l.c. \|\bar{z} \Lambda^{r+\frac{1}{2}} v\|^2$$

Now integrating by parts and commuting L_m and $\overline{L_m}$ we have

$$\|L_m \Lambda^{r-\frac{1}{2}} v\|^2 \leq \|\overline{L_m} \Lambda^{r-\frac{1}{2}} v\|^2 + \|\Lambda^r v\|^2$$

so that equation (2.1) becomes

$$(2.2) \quad \|\Lambda^r v\|^2 \lesssim \|\overline{L_m} v\|^2 + \|\overline{z} \Lambda^{r+\frac{1}{2}} v\|^2 \lesssim |(P_m v, v)| + \|\overline{z} \Lambda^{r+\frac{1}{2}} v\|^2$$

In the last term we integrate by parts, and obtain

$$\begin{aligned} \|\overline{z} \Lambda^{r+\frac{1}{2}} v\|^2 &= (\overline{z} \Lambda^{r+\frac{1}{2}} v, \overline{z} \Lambda^{r+\frac{1}{2}} v) = (z \overline{z} \Lambda^{r+1} v, \Lambda^r v) \\ &\leq s.c. \|\Lambda^r v\|^2 + \ell.c. \|z^2 \Lambda^{r+1} v\|^2 \end{aligned}$$

so that, iterating and absorbing the first term on the left in (2.2),

$$(2.3) \quad \|\Lambda^r v\|^2 \lesssim |(P_m v, v)| + \|z^{k-1} \Lambda^{r+\frac{k-1}{2}} v\|^2$$

To finish the derivation of the *a priori* estimate, we write $kz^{k-1} = [L_m, z^k]$, so that

$$\begin{aligned} (2.4) \quad k^2 \|z^{k-1} w\|^2 &= ([L_m, z^k] w, z^{k-1} w) \\ &= (L_m z^k w, z^{k-1} w) - (z^k L_m w, z^{k-1} w) = \tilde{A} + \tilde{B}. \end{aligned}$$

Then

$$|\tilde{A}| \lesssim |(z^{k-1} w, \overline{z} \overline{L_m} z^{k-1} w)| \lesssim s.c. \|z^{k-1} w\|^2 + l.c. \|\overline{L_m} w\|^2$$

since $[\overline{L_m}, z^{k-1}] = 0$, while

$$|\tilde{B}| \sim s.c. \|z^{k-1} w\|^2 + l.c. \|z^k L_m w\|^2.$$

The terms with the small constant (*s.c.*) will be absorbed on the left hand side of (2.4), yielding:

$$(2.5) \quad \|z^{k-1} w\|^2 \leq C \{ \|\overline{L_m} w\|^2 + \|\overline{z}^k L_m w\|^2 \}$$

Combining this estimate with (2.1), since $\|z^k w\| = \|\overline{z}^k w\|$, leads to

$$(2.6) \quad \|\Lambda^{-\frac{k-1}{2}} v\|^2 \leq C \{ \|\overline{L_m} v\|^2 + \|\overline{z}^k L_m v\|^2 \} = C |(P_m v, v)|.$$

with no errors (except the ‘microlocalizing ones, i.e., staying out of the elliptic region and the region z not equal to 0, where P_m has symplectic characteristic variety and [8], [10], and [9] apply).

Remark 1. *If one wishes to use the full pseudodifferential operator with symbol $(1 + |\zeta|^2 + |\tau|^2)^{1/2}$ instead of $\Lambda^{1/2}$, the derivation is virtually identical, with the addition of a lower order norm on the right of (2.6).*

3. THE LOCALIZATION OF POWERS OF T AND ITS COMMUTATORS

To apply (2.6) we must replace v by a localization of powers of $\partial/\partial t$, or, for convenience, we will localize powers of

$$T = -\frac{2i}{m+1} \frac{\partial}{\partial t}$$

applied to the solution u . Note that

$$[L_m, \overline{L_m}] = (m+1)^2 |z|^{2m} T,$$

which will be of technical use later.

For now we treat the solution u as being smooth, and concentrate on the estimates, remarking that by multiplying the Fourier Transform $\tilde{u}(z, \bar{z}, \tau)$ of u in t by dilations of $\Psi(\tau)$ equal to zero for $|\tau| > 2$ but identically equal to one for $|\tau| < 1$, once we obtain uniform estimates (in ε) for derivatives of the inverse transform of $\Psi(\tau/\varepsilon)(\tilde{u})$ we may let $\varepsilon \rightarrow 0$ to obtain the same estimates for the same derivatives of u .

The localization in space may be taken independent of $z, \bar{z}$ since taking a product of two localizers, $\varphi_1(t)\varphi_2(z, \bar{z})$ in terms where no derivatives land on φ_2 it just survives to the left of $P_m = f$ while in terms with derivatives, those terms are supported in the region where z is different from zero, i.e., in regions where the result is known, as remarked above.

But even with φ depending on t alone, the localization must be done very carefully. For example, the first bracket with P_m which we encounter will contain $(L_m \varphi(t))T^p \sim \bar{z}|z|^{2m} \varphi' T^p$, which is problematic for any value of m .

In Kohn's work [6] ($m = 0$), the $\bar{z}$ (or z) in front of each derivative of φ is carefully followed, and shown to provide, after some work, a gain of $1/2$ derivative. Analyticity was not considered in that paper, nor does it seem likely that it could be shown by those methods.

Derridj and Tartakoff found in [4] that an entirely different approach, involving a delicately balanced localization of T^p , led to analyticity rather directly.

Here, to handle the case $m > 0$, which corresponds to a CR manifold of finite type, we find it simpler to use a somewhat differently balanced localization. And with this localization both C^ω and C^∞ hypoellipticity come together.

3.1. The case $m = 0$. For motivation, however, we first we give the definition in the case $m = 0$.

Definition 3.1. For any pair of non-negative integers, (p_1, p_2) , let

$$(T_0^{p_1, p_2})_\varphi = \sum_{\substack{a \leq p_1 \\ b \leq p_2}} \frac{\overline{L}_m^a \circ \overline{z}^a \circ T^{p_1-a} \circ \varphi^{(a+b)} \circ T^{p_2-b} \circ z^b \circ L_m^b}{a!b!},$$

where

$$\varphi^{(r)} = \left(-i \frac{\partial}{\partial t} \right)^r \varphi.$$

Note that the leading term (with $a + b = 0$) is merely $T^{p_1} \circ \varphi \circ T^{p_2}$ which is equal to the operator $T^{p_1+p_2}$ on any open set Ω_0 where $\varphi \equiv 1$.

We have the extraordinary commutation relations:

Proposition 3.1.

$$[L_0, (T_0^{p_1, p_2})_\varphi] \equiv (T_0^{p_1, p_2-1})_{\varphi'} \circ L_0,$$

$$[\overline{L}_0, (T_0^{p_1, p_2})_\varphi] \equiv \overline{L}_0 \circ (T_0^{p_1-1, p_2})_{\varphi'},$$

$$[(T_0^{p_1, p_2})_\varphi, z] = z \circ (T_0^{p_1, p_2-1})_{\varphi'},$$

and

$$[(T_0^{p_1, p_2})_\varphi, \overline{z}] = (T_0^{p_1-1, p_2})_{\varphi'} \circ \overline{z},$$

where the $\equiv$ denotes modulo $C^{p_1-p'_1+p_2-p'_2}$ terms of the form

$$\frac{\overline{L}_0^{p_1-p'_1} \circ \overline{z}^{p_1-p'_1} \circ T^{p'_1} \circ \varphi^{(p'_1+p'_2+1)} \circ T^{p'_2} \circ z^{p_2-p'_2} \circ L_0^{p_2-p'_2}}{(p_1-p'_1)!(p_2-p'_2)!}$$

with either $p'_1 = 0$ or $p'_2 = 0$, i.e., terms where all free T derivatives have been eliminated on one side of φ or the other.

Proof. The proof is a straightforward calculation involving a shift of index in the definition of $(T_0^{p_1, p_2})_\varphi$. $\square$

3.2. The case $m > 0$. For $m > 0$, we use a related, but somewhat different definition of the localization, owing to multiple brackets. Parts of this discussion are in the style of [3], [2].

Definition 3.2. For $m > 0$, and (p_1, p_2) as above, set

$$\varphi^{(d)} = \left(i \frac{\partial}{\partial t} \right)^d \varphi(t),$$

$$N_b = \sum_{b' \leq q_b} A_{b'}^b \frac{(zL_m)^{b'}}{b'!},$$

where the $A_{b'}^b$ (real) are to be determined subject to $A_b^b = 1$, and

$$\tilde{N}_a = \sum_{a' \leq a} A_{a'}^a \frac{(-\overline{L_m} \bar{z})^{a'}}{a'!} = (N_a)^*$$

with the same coefficients, and finally we set

$$(T_m^{p_1, p_2})_\varphi = \sum_{\substack{a \leq p_1 \\ b \leq p_2}} \tilde{N}_a \circ T^{p_1-a} \circ \varphi^{(a+b)} \circ T^{p_2-b} \circ N_b.$$

We have

$$\begin{aligned} (3.1) \quad [\overline{L_m}, (T_m^{p_1, p_2})_\varphi] &= [\overline{L_m}, \sum_{a \leq p_1, b \leq p_2} \tilde{N}_a T^{p_1-a} \varphi^{(a+b)} T^{p_2-b} N_b] \\ &= \sum_{a \leq p_1, b \leq p_2} \left\{ [\overline{L_m}, \tilde{N}_a] T^{p_1-a} \varphi^{(a+b)} T^{p_2-b} N_b \right. \\ &\quad \left. - \tilde{N}_a T^{p_1-a} z |z|^{2m} \varphi^{(a+b+1)} T^{p_2-b} N_b + \tilde{N}_a T^{p_1-a} \varphi^{(a+b)} T^{p_2-b} [\overline{L_m}, N_b] \right\}. \end{aligned}$$

The last two terms on the right must cancel, to preserve the balance, since both disturb the balance between derivatives on φ and gain in powers of T . We will choose the coefficients $A_{b'}^b$ of N_b in such a way that, modulo acceptable errors,

$$(3.2) \quad [\overline{L_m}, N_b] = z |z|^{2m} T N_{b-1}.$$

This will provide the needed cancellation via a shift of index in a in the sum just as in the case with $m = 0$. The corresponding relation for brackets with L_m will follow by taking adjoints: again modulo acceptable errors,

$$(3.3) \quad [L_m, \tilde{N}_a] = -\tilde{N}_{a-1} \bar{z} |z|^{2m} T.$$

Condition (3.2) reads, using the definition of N_b , reads:

$$\sum_{b'=0}^b A_{b'}^b \frac{1}{b'!} [\overline{L_m}, (z L_m)^{b'}] = -\bar{z} |z|^{2m} T \sum_{b'=0}^{b-1} A_{b'}^{b-1} \frac{(z L_m)^{b'}}{b'!}.$$

Expanding the brackets and keeping all factors of $z L_m$ to the right,

$$\begin{aligned} \frac{1}{b'!} [\overline{L_m}, (z L_m)^{b'}] &= \frac{1}{b'!} \sum_{1 \leq b'' \leq b'} \binom{b'}{b''} a d_{z L_m}^{b''} (\overline{L_m}) (z L_m)^{b'-b''} \\ &= \frac{1}{b'!} \sum_{1 \leq b'' \leq b'} \binom{b'}{b''} a d_{z L_m}^{b''-1} ((m+1)^2 z |z|^{2m} T) (z L_m)^{b'-b''} \end{aligned}$$

$$= (z|z|^{2m}T) \sum_{1 \leq b'' \leq b'} \frac{(m+1)^{b''+1}}{b''!} \frac{(zL_m)^{b'-b''}}{(b'-b'')!}.$$

The condition (3.2) thus requires, renaming $b' - b''$ as $\bar{b}$ on the right just above,

$$(3.4) \quad \sum_{b''=1}^{b-\bar{b}} A_{b+b''}^b \frac{(m+1)^{b''+1}}{b''!} = A_{\bar{b}}^{b-1}.$$

Fortunately, we have investigated these equations in [3] and, citing a result in the book by Hirzebruch [5] have explicit solutions $\tilde{A}_*^*$ in the case that $m = 0$, unique under the conditions that $\tilde{A}_0^q = (-1)^q$, namely

$$\tilde{A}_s^r = \left(\left(\frac{t}{e^t - 1} \right)^{r+1} \right)^{(r-s)} (0)/(r-s)!$$

and for positive m we merely dilate by the factor of $m+1$:

$$A_s^r = \left(\left(\frac{\frac{t}{m+1}}{e^{\frac{t}{m+1}} - 1} \right)^{r+1} \right)^{(r-s)} (0)/(r-s)!$$

In addition, we will also need good expressions for the other brackets: we compute

$$(3.5) \quad \begin{aligned} [L_m, N_b] &= [L_m, \sum_{b'=0}^b A_{b'}^b \frac{(zL_m)^{b'}}{b'!}] = \sum_{\substack{1 \leq b'' \leq b' \\ b' \leq b}} A_{b'}^b \frac{1}{b''!} \frac{(zL)^{b'-b''}}{(b'-b'')!} \circ L_m \\ [N_b, z] &= z \circ \sum_{\substack{1 \leq b'' \leq b' \\ b' \leq b}} A_{b'}^b \frac{1}{b''!} \frac{(zL)^{b'-b''}}{(b'-b'')!} \\ [\overline{L_m}, \tilde{N}_a] &= -\overline{L_m} \circ \sum_{\substack{1 \leq a'' \leq a' \\ a' \leq a}} A_{a'}^a \frac{1}{a''!} \frac{(-\overline{L_m} \tilde{z})^{a'-a''}}{(a'-a'')!}, \\ [\tilde{z}, \tilde{N}_a] &= \sum_{\substack{1 \leq a'' \leq a' \\ a' \leq a}} A_{a'}^a \frac{1}{a''!} \frac{(-\overline{L_m} \tilde{z})^{a'-a''}}{(a'-a'')!} \circ \tilde{z}. \end{aligned}$$

In order to recognize these sums as N 's or $\tilde{N}$'s, we need to be able to shift the lower indices on $A_{a'}^a$ down by one. But this also we have done in [3], with the result that

Proposition 3.2. *For any r, s , and c ,*

$$A_s^r = \sum_{j=0}^{r-s} S_j^{r-s} A_{s-c}^{r-(c+j)}$$

where

$$|S_\ell^k| \leq C^k.$$

These brackets, then, together with the Proposition, immediately translate, setting $b'' = c$ and $\tilde{b} = b' - b''$, into:

$$[L_m, N_b] = \sum_{\tilde{b} \leq b-c} \frac{1}{c!} A_{b+c}^b \frac{(zL)^{\tilde{b}}}{\tilde{b}!} \circ L_m = \sum_{\tilde{b} \leq b-c-j} \frac{1}{c!} S_j^{c+j} A_{\tilde{b}}^{b-c-j} \frac{(zL)^{\tilde{b}}}{\tilde{b}!} \circ L_m$$

or

$$[L_m, N_b] = \sum_{\substack{c+j \leq b \\ 1 \leq c}} \frac{1}{c!} S_j^{c+j} N_{b-c-j} \circ L_m$$

Similarly,

$$[\overline{L_m}, \tilde{N}_a] = -\overline{L_m} \circ \sum_{\substack{c+j \leq a \\ 1 \leq c}} \frac{1}{c!} S_j^{c+j} \tilde{N}_{a-c-j},$$

and

$$[\overline{z}, \tilde{N}_a] = \sum_{\substack{c+j \leq a \\ 1 \leq c}} \frac{1}{c!} S_j^{c+j} \tilde{N}_{a-c-j} \circ \overline{z},$$

and

$$[N_b, z] = z \circ \sum_{\substack{c+j \leq b \\ 1 \leq c}} \frac{1}{c!} S_j^{c+j} \tilde{N}_{b-c-j},$$

These wonderful commutation relations mean that the whole localization $(T^{p_1, p_2})_\varphi$ may be commuted meaningfully with the vector fields $L_m, \overline{L_m}$ and with $z, \overline{z}$:

Proposition 3.3. *Modulo terms in which either p_1 or p_2 has been reduced to zero, and in view of the cancellations ensured by (3.2),*

$$(3.6) \quad [L_m, (T_m^{p_1, p_2})_\varphi] \equiv \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_2}} \frac{1}{c!} S_j^{c+j} (T_m^{p_1, p_2-(c+j)})_{\varphi^{(c+j)}} \circ L_m$$

Proof.

$$[L_m, (T_m^{p_1, p_2})_\varphi] \equiv \sum_{\substack{a \leq p_1 \\ b \leq p_2}} \tilde{N}_a \circ T^{p_1-a} \circ \varphi^{(a+b)} \circ T^{p_2-b} [L_m, N_b]$$

$$\begin{aligned}
&\equiv \sum_{\substack{a \leq p_1 \\ b \leq p_2}} \sum_{\substack{c+j \leq b \\ 1 \leq c}} \frac{1}{c!} S_j^{c+j} \tilde{N}_a \circ T^{p_1-a} \circ \varphi^{(a+b)} \circ T^{p_2-b} N_{b-c-j} \circ L_m \\
&\equiv \sum_{\substack{a \leq p_1 \\ b \leq p_2}} \sum_{\substack{c+j \leq b \\ 1 \leq c}} \frac{1}{c!} S_j^{c+j} \tilde{N}_a T^{p_1-a} \varphi^{(c+j)(a+b-c-j)} T^{p_2-(c+j)-(b-c-j)} N_{b-c-j} \circ L_m \\
&\equiv \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_2}} \frac{1}{c!} S_j^{c+j} (T_m^{p_1, p_2-(c+j)})_{\varphi^{(c+j)}} \circ L_m
\end{aligned}$$

□

Similarly we state, and omit the proofs, which are virtually identical to that of the previous proposition,

Proposition 3.4.

$$(3.7) \quad [\overline{L}_m, (T_m^{p_1, p_2})_\varphi] \equiv -\overline{L}_m \circ \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_1}} \frac{1}{c!} S_j^{c+j} (T_m^{p_1-(c+j), p_2})_{\varphi^{(c+j)}},$$

$$(3.8) \quad [\overline{z}, (T_m^{p_1, p_2})_\varphi] \equiv \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_1}} \frac{1}{c!} S_j^{c+j} (T_m^{p_1-(c+j), p_1})_{\varphi^{(c+j)}} \circ \overline{z},$$

and

$$(3.9) \quad [z, (T_m^{p_1, p_2})_\varphi] \equiv - \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_2}} \frac{1}{c!} S_j^{c+j} (T_m^{p_1, p_2-(c+j)})_{\varphi^{(c+j)}} \circ z$$

What these commutation relations mean is that we may move the vector fields of P_m past $(T_m^{p_1, p_2})_\varphi$ freely, at each stage incurring errors with the same vector fields and a gain in derivatives in $(T_m^{p_1, p_2})_\varphi$. Thus we may iterate the *a priori* inequality modulo errors of nearly arbitrarily low order - all of the $\equiv$ signs above mean that we will ultimately arrive at errors where either $p_1 = 0$ or $p_2 = 0$.

So we insert first $v = (T_m^{p_1, p_2})_\varphi u$ into (2.6), then bring $(T_m^{p_1, p_2})_\varphi$ to the left of $P_m = L\overline{L} + \overline{L}z^k \overline{z}^k L$, now writing L instead of L_m , since the formal expansions of the brackets are insensitive to m . so that we have:

$$\begin{aligned}
&\|\overline{L}(T_m^{p_1, p_2})_\varphi u\|_0^2 + \|\overline{z}^k L(T_m^{p_1, p_2})_\varphi u\|_0^2 + \|\Lambda^{-\frac{k-1}{2}} (T_m^{\frac{p}{2}, \frac{p}{2}})_\varphi u\|_0^2 \\
(3.10) \quad &\lesssim |(P_m(T_m^{p_1, p_2})_\varphi u, (T_m^{p_1, p_2})_\varphi u)_{L^2}| \\
&\lesssim |((T_m^{p_1, p_2})_\varphi P_m u, (T_m^{p_1, p_2})_\varphi u)_{L^2}| + |([P_m, (T_m^{p_1, p_2})_\varphi] u, (T_m^{p_1, p_2})_\varphi u)_{L^2}|
\end{aligned}$$

and by the above bracket relations, modulo the same terms as above where all T 's from one side of φ or the other have been 'converted' into L 's or $\bar{L}$'s, we have

$$\begin{aligned}
& ([P_m, (T_m^{p_1, p_2})_\varphi]u, (T_m^{p_1, p_2})_\varphi u) \equiv \\
& = ([L\bar{L}, (T^{p_1, p_2})_\varphi]u, (T^{p_1, p_2})_\varphi u) + ([\bar{L}z^k\bar{z}^k L, (T^{p_1, p_2})_\varphi]u, (T^{p_1, p_2})_\varphi u) \\
& = ([L, (T^{p_1, p_2})_\varphi]\bar{L}u, (T^{p_1, p_2})_\varphi u) + (L[\bar{L}, (T^{p_1, p_2})_\varphi]u, (T^{p_1, p_2})_\varphi u) \\
& + ([\bar{L}, (T^{p_1, p_2})_\varphi]z^k\bar{z}^k Lu, (T^{p_1, p_2})_\varphi u) + (\bar{L}[z^k, (T^{p_1, p_2})_\varphi]\bar{z}^k Lu, (T^{p_1, p_2})_\varphi u) \\
& + (\bar{L}z^k[\bar{z}^k, (T^{p_1, p_2})_\varphi]Lu, (T^{p_1, p_2})_\varphi u) + (\bar{L}z^k\bar{z}^k[L, (T^{p_1, p_2})_\varphi]u, (T^{p_1, p_2})_\varphi u) \\
& \equiv \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_2}} \frac{1}{c!} S_j^{c+j} ((T_m^{p_1, p_2-(c+j)})_{\varphi(c+j)} L\bar{L}u, (T_m^{p_1, p_2})_\varphi u) \\
& - \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_1}} \frac{1}{c!} S_j^{c+j} (L\bar{L}(T_m^{p_1-(c+j), p_2})_{\varphi(c+j)} u, (T_m^{p_1, p_2})_\varphi u) \\
& - \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_1}} \frac{1}{c!} S_j^{c+j} (\bar{L}(T_m^{p_1-(c+j), p_2})_{\varphi(c+j)} z^k\bar{z}^k Lu, (T_m^{p_1, p_2})_\varphi u) \\
(3.11) \quad & - \sum_{k'=1}^k \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_2}} \frac{1}{c!} S_j^{c+j} (\bar{L}z^{k'}(T_m^{p_1, p_2-(c+j)})_{\varphi(c+j)} z^{k-k'}\bar{z}^k Lu, (T_m^{p_1, p_2})_\varphi u) \\
& + \sum_{k'=0}^{k-1} \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_1}} \frac{1}{c!} S_j^{c+j} (\bar{L}z^k\bar{z}^{k'}(T_m^{p_1-(c+j), p_1})_{\varphi(c+j)} \bar{z}^{k-k'} Lu, (T_m^{p_1, p_2})_\varphi u) \\
& + \sum_{\substack{1 \leq c, 0 \leq j \\ c+j \leq p_2}} \frac{1}{c!} S_j^{c+j} (\bar{L}z^k\bar{z}^k (T_m^{p_1, p_2-(c+j)})_{\varphi(c+j)} Lu, (T_m^{p_1, p_2})_\varphi u) \\
& = A_1 + A_2 + A_3 + \sum_{k'=1}^k A_{4, k'} + \sum_{k'=0}^{k-1} A_{5, k'} + A_6.
\end{aligned}$$

Concerning the critical $L, \bar{L}, z^k$ and $\bar{z}^k$, note that in each term above

- no $L, \bar{L}$, power of z or power of $\bar{z}$ has been lost,
- the order among $L, \bar{z}^k, z^k$, and $\bar{L}$ is preserved,
- letting $|q| = q_1 + q_2$, each term on the right contains $(T^{q_1, q_2})_{\varphi(|p|-|q|)}$ with $|q| < |p|$ and $|p| - |q|$ derivatives on φ ,

- just as (3.11) demonstrates the errors which result in moving $(T^{p_1, p_2})_\varphi$ past the vector fields $L, \overline{L}_m, z^k \overline{L}_m$ and $\overline{z}^k L$, further such brackets to position the vector fields so as to make use the *a priori* estimate again will produce similar errors, with $|q|$ still lower and the ‘lost’ T derivatives transferred to φ ,
- iterating this process, together with a weighted Schwarz inequality, will produce a sum of terms with $q_j \leq \frac{p}{2}$ of the form

$$l.c. \|\Lambda^{\frac{k-1}{2}} (T_m^{q_1, q_2})_{\varphi(|p|-|q|)} P u\|_0^2 + s.c. \|\Lambda^{-\frac{k-1}{2}} (T_m^{\frac{p}{2}, \frac{p}{2}})_\varphi u\|^2.$$

- In fact, using larger constants $\tilde{S}_j^{c+j}$ subject to the same kind of bounds, $|\tilde{S}_j^{c+j}| \leq \tilde{C}^{c+j}$, we may replace all sums on the right hand side above by suprema subject to the same range restrictions on the indices.
- This use of suprema allows us easily to iterate everything on the right with easy control on the constants until either p_1 or p_2 , both of which start as $\frac{p}{2}$, drops to zero, which may happen in two ways - either by stepwise decrease as on the right hand side above from successive brackets or by the single term in Propositions 3.1 and 3.3 which is not cancelled, the term with all L ’s or $\overline{L}$ ’s on one side or the other in the definition of $(T_m^{p_1, p_2})_\varphi$, whose principal term is $(T_m^{p_1, 0})_{\varphi(p_2)} (zL)^{p_2}/p_2!$ or its analogue with p_1 reduced to 0.
- At this point we no longer have an effective localization of powers of T - for example, brackets with $\overline{L}$ are not corrected. We proceed anyway, and when we lack a ‘good’ vector field such as $\overline{L}$ (or of course $\overline{z}^k L$), we create one by integrating by parts:

$$(3.12) \quad \|Lw\|^2 \leq \|\overline{L}w\|^2 + |(|z|^{2m})Tw, w|$$

to use up the L and $\overline{L}$ derivatives with the byproduct of introducing up to half the number of new T derivatives.

Overall, then, the strategy has been, with universal C , bounding derivatives of the Ehrenpreis-type localizing functions $\varphi^{(r)}$ by $(CN)^r \sim C^r r!$ for $r \leq N \sim p$:

$$(3.13) \quad \frac{\|\Lambda^{-\frac{k-1}{2}} T^p u\|_{\{\varphi \equiv 1\}}}{N^p} \rightarrow \frac{\|\Lambda^{-\frac{k-1}{2}} (T_m^{\frac{p}{2}, \frac{p}{2}})_\varphi u\|}{N^p} \rightarrow \\ \rightarrow C^{\frac{p}{4}} \frac{\|\Lambda^{-\frac{k-1}{2}} T^{\frac{3p}{4}} u\|_{\text{supp } \varphi}}{N^{\frac{3p}{4}}} \rightarrow C^{\frac{p}{4}} \frac{\|\Lambda^{-\frac{k-1}{2}} T^{\frac{3p}{4}} u\|_{\{\varphi_1 \equiv 1\}}}{N^{\frac{3p}{4}}} \rightarrow \dots$$

for suitable $\varphi_1 \equiv 1$ on the support of φ . This will continue, with a sequence of φ_j supported in nested intervals as in [8], [9] until only a

negligible fraction of p is left, namely a bounded number of derivatives. Since the order in T^p is reduced by a factor of $3/4$ each time, we will need $\log_{4/3} p$ such nested open sets. Thus:

$$(3.14) \quad \frac{\|\Lambda^{-\frac{k-1}{2}} T^p u\|_{\{\varphi \equiv 1\}}}{N^p} \lesssim C^p \|\Lambda^{-\frac{k-1}{2}+3} u\|_{\{\varphi_{\log_{4/3} p} \equiv 1\}}$$

where the 3 could be any other small integer. And of course the whole derivation could have been done at the H^s level: for any given s ,

$$(3.15) \quad \frac{\|\Lambda^{s-\frac{k-1}{2}} T^p u\|_{\{\varphi \equiv 1\}}}{N^p} \lesssim C^p \|\Lambda^{s-\frac{k-1}{2}+3} u\|_{\{\varphi_{\log_{4/3} p} \equiv 1\}}$$

which will end the story if this last norm is known to be finite even as the localizer in τ tends to the identity, provided that the terms that arise along the way are all similarly bounded. The most important of these is of course

$$|(\Lambda^{s+\frac{k-1}{2}} (T_m^{\frac{p}{2}, \frac{p}{2}})_\varphi P u, \Lambda^{s-\frac{k-1}{2}} (T_m^{\frac{p}{2}, \frac{p}{2}})_\varphi u)|$$

which shows that $P_m u \in H^{s+\frac{k-1}{2}+p}$ implies that $u \in H^{s-\frac{k-1}{2}+p}$, a loss of $k-1$ derivatives.

Remark 1. *The value of s will be chosen so that we know the norm on the right in (3.15) is finite as the localizer in τ goes to the identity, and then p will be chosen so that $P_m u \in H^{s+\frac{k-1}{2}+p}$ for that value of s . It follows that $u \in H^{s-\frac{k-1}{2}+p}$.*

Remark 2. *For analyticity, one needs to ensure that as we take p larger and larger, the constants satisfied by the Ehrenpreis-type localizers are subject to bounds such that the estimate (3.15) is uniform in p . We have shown this often before and the arguments are the same here.*

REFERENCES

- [1] A. BOVE, M. DERRIDJ, J.J. KOHN AND D.S. TARTAKOFF, *Hypoellipticity for a sum of squares of complex vector fields with large loss of derivatives*, preprint.
- [2] A. BOVE AND D.S. TARTAKOFF, *Analytic hypo-ellipticity at non-symplectic characteristics when the symplectic form changes its rank*, preprint.
- [3] M. DERRIDJ AND D.S. TARTAKOFF, *Local analyticity for $\square_b$ and the $\bar{\partial}$ -Neumann problem at certain weakly pseudoconvex points*, Commun.in Partial Differential Equations **13(12)** (1988), pp. 1521-1600.
- [4] M. DERRIDJ AND D.S. TARTAKOFF, *Analyticity and loss of derivatives*, preprint, submitted to Annals of Mathematics.
- [5] F. HIRZEBRUCH, *Topological Methods in Algebraic Geometry*, Springer Verlag, New York, 1966.

- [6] J.J. KOHN, *Hypoellipticity and loss of derivatives*, Annals of Mathematics, to appear.
- [7] C. PARENTI AND A. PARMEGGIANI, *A Note on Kohn's and Christ's Examples*, preprint.
- [8] D.S. TARTAKOFF, *Local Analytic Hypoellipticity for $\square_b$ on Non-Degenerate Cauchy Riemann Manifolds*, Proc. Nat. Acad. Sci. U.S.A. **75** (1978), pp. 3027-3028.
- [9] D.S. TARTAKOFF, *On the Local Real Analyticity of Solutions to $\square_b$ and the $\bar{\partial}$ -Neumann Problem*, Acta Math. **145** (1980), pp. 117-204.
- [10] F. TREVES, *Analytic Hypo-ellipticity of a Class of Pseudo-Differential Operators with Double Characteristics and Application to the $\bar{\partial}$ -Neumann Problem*, Comm. in P.D.E. **3** (6-7) (1978), pp. 475-642.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO, M/C
249, 851 S. MORGAN ST., CHICAGO IL 60607, USA
E-mail address: `dst@uic.edu`